

WEATHER DATA FOR THE PREVAILING WEEK
Date of Foundation Pruning: 15/04/2021
Wednesday (1/09/2021)–Wednesday (08/09/2021)

Location	Temperature (°C)		Possibility of Rain	Cloud Cover	Wind Speed (Km/hr) Min-Max	R H%	
	Min	Max				Min	Max
Nashik	22-23	27-30	Nashik, Dindori, Ozar, Palkhed, , Vani, Loni, Pimpalgaon Baswant, Shirdi, Kalwan Wed- Good Rain. Thu to Mon- Light Rain.	Mostly Cloudy	15-22	71-86	94-96
Pune	20-21	26-27	Pune, Phursungi, Loni Kalbhor, Uruli Kanchan, Patas, Yavat, Supa, Narayangaon, Baramati Wed- Light to Moderate Rain.	Partly Cloudy	17-20	59-67	87-90
Solapur	20-21	29-32	Solapur, Vairag, Nannaj, Kati, Pangri, Osmanabad, Latur, Ausa, Tuljapur, Pandharpur, Barshi, Atpadi, Kasegaon Wed to Sat- Light to Moderate Rain. Sun to Tue- Good Rain.	Mostly Cloudy	15-19	51-62	82-89
Sangli	21	27-29	Sangli, Miraj, Palus, Kagvad, Shetfal, Palsi, Khanapur, Vita, Tasgaon, Shirguppi, Arag, Walva, Kawthe Wed, Thu & Sat- Light Rain. Fri- Good Rain. Sun to Tue- Moderate Rain.	Mostly Cloudy	14-19	63-71	92-95
Vijayapura	20-21	28-31	Vijayapura, Chadchan, Tikota, Telsang Thu to Tue- Moderate to Good Rain.	Mostly Cloudy	21-24	57-66	86-89
Hyderabad	22	29	Hyderabad, Medchal, Zahirabad Thu to Next Wed- Good Rain.	Mostly Cloudy	11-18	62-73	87-93
Satara	20-21	25-27	Satara, Phaltan, Man, Khatav Rahata Thu to Tue- Light to Moderate Rain.	Partly to Mostly Cloudy	10-14	65-72	92-94
Ahmednagar	21-22	28-31	Ahmednagar, Nagar, Kopargaon, Shrigonda, Karjat, Jamkhed, Akole, Rahata, Sangamner Sat- Drizzling. Sun to Tue- Good Rain.	Mostly Cloudy	12-23	58-64	86-91
Jalna	22-23	30-31	Jalna, , Jafrabad, Mantha, Ambad, Gansawangi Sat to Next Wed- Good Rain.	Mostly Cloudy	12-17	61-64	86-92
Buldhana	23-24	30-32	Buldana, Chikhli, D.raja, Sindkhedraja Thu to Next Wed- Good Rain.	Mostly Cloudy	08-20	67-71	89-96
Kolhapur	23-24	28-29	Gagan-bavada, Kagal, Karveer Thu to Tue- Moderate to Good Rain.	Mostly Cloudy	07-09	76-80	97-98

Bengaluru Rural	18-20	26-29	Bangaluru-east, Bangaluru-north, Bangaluru-south, Doddaballapur, Anekal Thu to Next Wed- Good Rain.	Mostly cloudy	14-20	55-64	87-91
Belagavi	22-23	29	Belagavi, Athni, Chikodi, Gokak, Khanapur Fri & Tue- Moderate Rain. Sat- Light Rain.	Mostly cloudy	14-18	68-74	93-98
Bidar	20-21	28-29	Bidar, Basavakalyan, Humnabad Thu to Next Wed- Good Rain.	Mostly cloudy	12-15	63-72	91-94
Bagalkot	21	29-30	Bagalkot, Bilagi, Jamkhandi, Mudhol, Hungund, Badami Thu to Next Wed- Moderate to Good Rain.	Mostly cloudy	19-22	57-63	86-89

Note: Above weather information is summary of weather forecasting given in following websites

https://www.wunderground.com/?cm_ven=cgi

<https://imdagrimet.gov.in/weatherdata/BlockWindow.php>

<https://www.accuweather.com/>

ICAR-National Research Centre for Grapes does not claim accuracy of it.

II. Water management (Dr. A.K. Upadhyay)

a) Days after foundation pruning: 138

b) Pan evaporation: : Nil - 3mm

Amount of irrigation advised:

1. All the grape growing regions are forecasted to receive rains. The irrigation water application should be based upon the growth of the vines. Objective is to concentrate on cane maturity, hence, vigour should be controlled.
2. **Cane maturity stage:** As the irrigation water requirement is low and rains expected, donot irrigate the vineyards. If no rains for more than 3days and vines are showing stress, then, apply irrigation through surface drip @ 1500 to 2000 L/acre per day.
3. If continuous good rains are forecasted, remove the mulch and allow the bund/ rootzone to be fully wet with water for leaching of salts. The mulch so removed can be mixed with the soil to improve the soil porosity. This is especially important for the following condition:
 - i) In Solapur, Sangli, Vijayapura or any area where the ground water used for irrigation contains more salt.

Nutrient management

1. After current rains, give foliar spray of SOP @ 3-5 g/L depending upon canopy.
2. In case of calcareous soils where acute iron deficiency is observed, repeatedly spray 2-3g/L Ferrous sulphate two to three times at 3 days interval followed by 15-20 kg/ acre Ferrous sulphate application through drip. The fertigation dose should be split into atleast 3 doses of 5kg each. Apply 5kg/ acre soluble sulphur through drip every week. Also spray magnesium sulphate and potassium sulphate @ 3 gm each/ L once only. Keep a close watch on the development of leaf blackening symptoms if irrigation water contains sodium more than 100ppm.
3. Possibility of leaf curling, check the leaf margins, if slight to more yellow, possibility of potassium deficiency. Foliar spray of SOP @ 3-4g/L followed by fertigation of 20-25 kg SOP/acre in 2 to 3 splits.
4. If the leaf yellowing starts from in between the leaf veins then, possibility of magnesium deficiency is there. Foliar spray of Magnesium sulphate @ 3-4g/L followed by fertigation of 15-20 kg magnesium sulphate/acre in 2 to 3 splits.
5. In coloured varieties like Jumbo, Nanasaheb Purple Seedless etc., leaf curling along with reddening/ bronzing of the leaf margin can be observed if potassium deficiency is there. Foliar spray of SOP @ 3g/L followed by fertigation of 20-25 kg SOP/acre in 2 to 3 splits.
6. In calcareous soils, provide foliar application of Sulphate of Potash and Magnesium sulphate each (@ 4g/L once in this growth stage).

Pre-pruning operations – Fruit pruning season

1. In case pruning is planned during August - September, raise Sunnhemp or Dhaincha for green manuring purpose.
2. The vineyards where sodicity problems are there, apply gypsum to the soil for removal of sodium from the soil exchange complex. In case of calcareous soils, use sulphur for similar purpose. The application should be alongwith FYM/compost etc. They should be mixed in the soil and not left on the top.
3. In case of calcareous soils, if SSP is applied as basal dose, mix with FYM/compost etc. to avoid phosphorus fixation.
4. Test the soil and irrigation water, to plan for nutrient and water management during fruit pruning season.

5. In areas where rains have not been received and the irrigation water availability is less, it is suggested to flood the rootzone(only) with water to leach out the salts and wet the entire soil depth before pruning and then cover with mulch. Thereafter irrigate as per availability of water.

III. Requirement of growth regulators (Dr. S.D. Ramteke)

Nil

IV. Canopy management (Dr. R.G. Somkuwar)

Grafting to be done:

1. Selection of rootstock:

2. The shoot selected for grafting should be either semi matured or soft wood.
3. The rootstock shoot should be straight growing, healthy and free from disease
4. The rootstock shoot should be in complete sap flow condition.
5. The diameter of rootstock should be 8-10mm at 1 feet above the ground surface.
6. Under the condition of less rainfall, the rootstock plants need to be irrigated 3-4 days before actual grafting. This will help to form sap in the rootstock plant.

7. Selection of scion:

8. The scion selected from the vine should be free from diseases.
9. The scion selected for grafting should be regular bearer, high yielding and true to type
10. The scion selected should be completely matured contains food material.
11. The scion selected should be round in shape.
12. The scion should be dipped in Carbendazim solution for about 2-3 hours before grafting.

13. Condition for grafting success:

14. While performing the grafting and after, the temperature in the vineyard should be 30-35°C and the relative humidity should be above 80%.
15. The person performing the grafting should possess the skill.
16. The tape used for grafting should have proper elasticity.

V. Disease management (Dr. Sujoy Saha)

Days after foundation pruning	Risk of diseases			
	Downy mildew	Powdery mildew	Anthracnose	Others (specify)
138	Low	Low	High	Bacterial spot-High Rust- moderate

Rust had been observed in many orchards where there was water stagnation in the recent rains last week. Application of chlorothalonil @ 2g/l will control the disease. If the incidence is severe, application of flusilazole @ 25 mL/200L may be done, but it should not be applied at any stage after fruit pruning. Thiophenate methyl @1g/L may be given to protect from anthracnose disease. In areas of Nashik, Sangli, Osmanabad and Solapur, where bacterial spot is incident application of Mancozeb @2.5g/litre may be given. This will give an additional control of downy mildew. Application of Streptocycline in grapes is not advisable. Dusting of Mancozeb @4-5kg/acre during this wet and humid conditions will be effective against downy mildew. Drip application of Trichoderma may be continued in areas receiving rainfall. Foliar spray of Trichoderma may also be given @2- 3ml/L but it should not be given immediately after application of copper fungicides. If bacterial spot and anthracnose are incident together a ready-mix of kasugamycin + copper oxychloride @0.75g/l may be applied twice at an interval of 10 days. If fruit pruning is done in Satana and Indapur region, stem and cordon wash with mancozeb 75WP @2.5-3g/l followed by sulphur@ 2.5-3g/l should be done at an interval of 3-5 days.

VI. Insect and Mite management. (Dr. D.S. Yadav)

Days after pruning	Risk of pests				
	Mealybug	Mite	Thrips	Caterpillar	Flea beetle
Cane maturity and afterwards	Moderate	Moderate	Moderate	High	Low to moderate

1. Stem borer, *Celosterna scabrator* adults may seen in vineyards and/or near light at night at homes near vineyards. In case, fipronil 80 WG @ 0.0625 g per litre water is used for caterpillars, it will also control these adults of stem borer.
2. In case of caterpillar infestation, application of fipronil 80 WG @ 0.0625 g per litre or emamectin benzoate 5 SG @ 0.22 g per litre or cyantraniliprole 10 OD @ 0.7 ml per litre water is effective.
3. Use of broad-spectrum insecticides should be avoided for mealybug control. Buprofezin 25 SC @ 1.25 ml per litre or spirotetramat 15.31 OD @ 0.7 ml per litre water may be given to manage mealybugs. Preventive plant wash, on stem and cordons, of biocontrol agents such as *Verticillium*, *Metarhizium*, *Beauveria* may be given.
4. In case of thrips infestation, remove excess shoot growth.
5. Red colour stem borer (*Dervishiya cadambae*) has started egg laying and infestation under bark in grape areas. Install light traps near the vineyards to manage moths of this stem borer. Remove loose bark from stem and cordons and give preventive wash on stem and cordons with biocontrol agent *Metarhizium* @ 3-5 ml per litre water minimum once in the month during July to September months. If infestation is observed, remove the loose bark and give stem and cordon wash with lambda cyhalothrin 5 CS @ 2.5 ml per litre water and 1.5-2 litres water per plant.
6. In new vineyards after grafting, flea beetle infestation may be observed. In case of heavy infestation, give soil drenching with imidacloprid 17.8 SL @ 1.5 ml per plant and foliar application with spinosad 45 SC @ 0.25 ml per litre or spinetoram 11.7 SC @ 0.3 ml per litre or fipronil 80 WG @ 0.0625 g per litre water.
7. Mite infestation may start appearing, therefore, monitor the vineyards carefully. If mite infestation is observed, sulphur 80 WDG @ 1.5-2.0 gram per litre or abamectin 1.9 EC @ 0.75 ml/l water is effective.